arXiv:0709.0385v1 [nucl-th] 4 Sep 2007

Interpretation of recent JLab results on quasi elastic (e,e'p) reactions off few-nucleon systems

C. Ciofi degli Atti [a]

[a] *Department of Physics, University of Perugia and Istituto Nazionale di Fisica Nucleare, Sezione di Perugia, Via A. Pascoli, I-06123, Italy*

L. P. Kaptari [a,b]

[b] *Bogoliubov Lab. of Theoretical Physics, 141980, JINR, Dubna, Russia*

H. Morita

Sapporo Gakuin University, 11 Bunkyo-dai, Ebetsu, Hokkaido 069-8555, Japan

---

Abstract

Recent JLab experimental data on quasi elastic $^3He(e,e'p)^2H(pn)$ and $^4He(e,e'p)^3H$ processes are interpreted using an approach based upon realistic wave functions and Glauber multiple scattering theory within a generalized eikonal approximation (GEA). The results of a non factorized calculation of the left-right asymmetry A_{TL} of the process $^3He(e,e'p)^2H$, obtained using the full covariant form of the electromagnetic operator, are also presented.

Key words: Glauber theory, Generalized Eikonal approximation, Realistic wave function
PACS: 21.45.-v, 24.10.-i, 25.10.+s, 25.30.-c

---

1. Introduction

The exclusive process A(e,e'p)B in few-nucleon systems is one of the most promising tools to investigate proton propagation in the medium. We have analyzed this process on the basis of realistic wave functions and a generalized eikonal approximation [1,2] to describe the final state interaction. In this contribution we will present the results of our interpretation of recent JLab data on the quasi elastic reactions $^3He(e,e'p)^2H(pn)$ and $^4He(e,e'p)^3H$.

Email addresses: `ciofi@pg.infn.it` (C. Ciofi degli Atti), `kaptari@thsun1.jinr.ru` (L. P. Kaptari), `hiko@sgu.ac.jp` (H. Morita).

2. Formulation

Within the factorization approximation, the differential cross section for the quasi-elastic A(e,e'p)B reaction can be written as follows

$$\frac{d^6\sigma}{d\omega d\Omega_e dp d\Omega_p} = K\sigma_{ep} S_D(\mathbf{p}_m, E_m), \tag{1}$$

where K is a kinematical factor, σ_{ep} the electron-proton cross section and $S_D(\mathbf{p}_m, E_m)$ the nuclear spectral function distorted by the final state interaction (FSI), which is treated here within the generalized eikonal approximation (GEA)[1,2]. In the case of few-nucleon system $S_D(\mathbf{p}_m, E_m)$ can be calculated using realistic wave functions and experimentally determined nucleon-nucleon scattering amplitude [2,3,6]. The violation of the factorization approximation will be discussed in the next Section.

3. Results

The results for the ${}^3He(e,e'p)^2H(pn)$ reactions [2,3] are presented in Fig.1 and Fig.2 respectively; a very good agreement between the GEA calculation and the experimental data can be seen. Fig. 1 shows that the cross section exhibits different slopes, which corresponds to the PWIA and to the single- and double-rescattering contributions, respectively [3].



Fig. 1. Results for the ${}^3He(e,e'p)^2H$ reaction *vs* the missing momentum p_m [3]. Dots: PWIA ; dash: FSI (single rescattering); full: FSI (single plus double rescattering). Experimental data from [4].



Fig. 2. The same as in Fig. 1 but for the process ${}^3He(e,e'p)pn$ *vs* the excitation energy E_{rel} of the $n-p$ pair ($E_{th} = E_3$ is the two-nucleon emission threshold in ${}^3He$).

Our results [6] for the ${}^4He(e,e'p)^3H$ process are shown in Fig. 3, and again a satisfactory agreement with the experimental data can be seen.

Finally in Fig. 4 the left-right asymmetry A_{TL}



Fig. 3. The reduced cross section $n_D(\mathbf{p}_m) = [d^5\sigma/(d\nu d\Omega_e d\Omega_p)] \times [\mathcal{K}\sigma_{ep}]^{-1}$ for the process $^4He(e,e'p)^3H$ [6]. Preliminary experimental data from [5].



Fig. 4. The asymmetry A_{TL} for the process $^3He(e,e'p)^2H$ [7]. Dot-dash: PWIA; dashed and full: nonfactorized results corresponding to single- and double-rescattering FSI, respectively. Experimental data from [4]

.

$$A_{TL} = \frac{\sigma(\phi = 0^{\circ}) - \sigma(\phi = 180^{\circ})}{\sigma(\phi = 0^{\circ}) + \sigma(\phi = 180^{\circ})}, \tag{2}$$

calculated within a non-factorized momentum space approach ,without using any kind of non-relativistic reduction, is shown Fig. 4 [7]; it can be seen that the factorized calculation cannot describe the data at $p_m \geq 200(MeV/c)$, whereas the non-factorized calculation can reproduce the data up to $p_m \simeq 600(MeV/c)$, but fails at higher momenta. The extension of the non factorized approach to the process $^4He(e,e'p)^3H$ is in progress [8].

4. Summary

We have analyzed recent JLab experimental data on quasi-elastic processes off few-nucleon systems using realistic wave functions and the Generalized Eikonal Approximation to treat the FSI. The factorized calculations reproduces very well the three-body break up and the right wing of of the two-body break. Our non-factorized calculation of the asymmetry A_{TL} does very well up to $p_m \simeq 600(MeV/c)$, but a strong disagreement between theory and experiment at higher momenta, common to many calculations, remains to be explained.

References

[1] M.M. Sargsian, T.V. Abrahamyan, M.I. Strikman and L.L. Frankfurt, *Phys. Rev.* **C71**(2005)044614.

[2] C. Ciofi degli Atti and L.P. Kaptari, *Phys. Rev.* **C71**(2005)024005.

[3] C. Ciofi degli Atti and L.P. Kaptari, *Phys. Rev. Lett.* **95**(2005)052502.

[4] F. Benmokhtar et al., *Phys. Rev. Lett.* **94**(2005)082325 and M. M. Rvachev et al., *Phys. Rev. Lett.* **94**(2005)192302.

[5] B. Reitz et al., *Eur. Phys. J.A* **S19**(2004)165.

[6] C. Ciofi degli Atti, L.P. Kaptari and H. Morita, *Nucl. Phys.* **A782**(2007)191.

[7] C. Ciofi degli Atti and L.P. Kaptari, nucl-th/0705.3951.
[8] C. Ciofi degli Atti, L.P. Kaptari and H. Morita, to appear.